

On cages with given degree sets

Denis Hanson

University of Regina, Regina, Sask., Canada S4S 0A2

Ping Wang

University of Regina, Regina, Sask., Canada S4S 0A2

Leif K. Jørgensen

Aalborg University, Aalborg, Denmark

Received 24 December 1990

Revised 11 November 1991

Abstract

Hanson, D., P. Wang and L.K. Jørgensen, On cages with given degree sets, *Discrete Mathematics* 101 (1992) 109–114.

We consider the problem of constructing minimal graphs of given girth having a particular degree set.

1. Introduction

A (v, g) -cage (or cage) is a v -regular (simple) graph of girth g on the minimum number of vertices. For $v = 2$ the cage is a cycle; for $g = 3$ the cage is K_{v+1} ; and for $g = 4$ the cage is $K_{v,v}$. The most famous cage is of course the $(3, 5)$ -cage or the Petersen graph, Fig. 1 (see [4]).

The exact order of cages is a difficult and long standing problem. A comprehensive article in this area is the survey paper of Wong [5]. Cages have been generalized in a number of ways such as prescribing the minimum ‘odd and even girth’. Another such generalization is as follows: if $D = \{a_1, a_2, \dots, a_k\}$ is a set of positive integers with $2 \leq a_1 < \dots < a_k$ then $f(D; g) = f(a_1, a_2, \dots, a_k; g)$ is the minimum order of a graph with degree set D and girth g . Thus for example the ordinary (v, g) -cage has order $f(v; g)$. Few values of $f(D; g)$ are known beyond the special cases (see Chartrand et al [2]):

$$f(3, 4; 5) = 13 \text{ and } f(3, 4; 6) = 18;$$

$$f(D; 3) = a_k + 1;$$

$$f(2, m; g) = 1/2(m(g-2) + 4) \text{ for } g \text{ even and}$$

$$f(2, m; g) = 1/2(m(g-1) + 2) \text{ for } g \text{ odd; and}$$

$$f(r, s; 4) = r + s.$$




Fig. 1.

In [3], Downs et al determined $f(3, m; 5) = f(3, 4, m; 5) = 3m + 1$; $f(3, m; 7) = 7m + 1$; and $f(3, m; 9) = 15m + 1$. In this article we consider the problems of evaluating $f(3, m; 6)$ and $f(4, m; 5)$; in particular we conclude that $f(3, m; 6) = 4m + 2$ for $m \geq 3$ and that $f(4, m; 5) = 4m + 1$ for $m \geq 5$.

2. The $(3, m; 6)$ -cages

To evaluate $f(3, m; 6)$ we will first show that the lower bound is $4m + 2$ and then use a construction to attain this bound. Suppose on the contrary that $f(3, m; 6) < 4m + 2$ and that G is a $(3, m; 6)$ -cage. For convenience we choose a vertex x of degree m ; let its neighbourhood set be denoted by $X = \{x_1, x_2, \dots, x_m\}$; let the set of vertices at distance two from x be denoted by X' ; finally denote the remaining vertices of G by Y where $|Y| = k$. We define two subgraphs, U and V , of a graph to be d -remote if the distance between them is at least d . We claim that any two points of degree m in G are 3-remote. Suppose for example that x_1 has degree m . Then girth 6 implies that the remaining neighbours of x_1 are not adjacent to the neighbours of x_2, \dots, x_m and we have $f(3, m; 6) \geq 1 + m + (m - 1) + (2(m - 1)) + 2(m - 1) = 6m - 4 \geq 4m + 2$ whenever $m \geq 3$. Thus we may assume that $X' = \{x_{11}, x_{12}, x_{21}, x_{22}, \dots, x_{m1}, x_{m2}\}$ and $|Y| \leq m$. Now suppose for example that x_{11} has degree m , then x_{11} has $m - 1$ neighbours in




Fig. 2.




Fig. 3.

Y and x_{12} has two further neighbours in Y , a contradiction. The 3-remoteness of such vertices says that G has the structure shown in Fig. 2 where all degrees in X and X' are equal to three.

If Y contains t points of degree m where $0 \leq t \leq 1$ then there are at most $tm + 3(k - t) \leq (3 + t)m - 3t$ edges from Y to X' , however this must be $4m$ using the girth 6 condition on the vertices of X' , a contradiction. Suppose that $t \geq 3$; since these t vertices must be 3-remote we have that $|X' \cup Y| \geq 3m + 3$ and $f(3, m; 6) > 4m + 1$. Thus we must have precisely $t = 2$. Without loss of generality let y_1 and y_2 have degree m .

Let A be the subset of X consisting of those vertices at distance two from both y_1 and y_2 ; let B be the subset of X consisting of those vertices at distance two from exactly one of y_1 or y_2 ; let $C = X - (A \cup B)$. Let the neighbourhood set of A in X' be A' . Finally let Y_1 and Y_2 be the (disjoint) neighbourhood sets of y_1 and y_2 respectively in Y and let $Y_3 = Y - (Y_1 \cup Y_2 \cup \{y_1, y_2\})$. Note that no vertex y in $Y_1 \cup Y_2$ is at distance two from A else G would contain a 3-cycle or a 5-cycle. Similarly no vertex in Y_3 can have three neighbours in A' for this would force a 4-cycle with either y_1 or y_2 . These two observations imply that each vertex in A' has at least one neighbour in Y_3 , and that $|Y_3| \geq |A|$. Let E^+ denote the set of edges joining y_1 and y_2 to X' . Then $|E^+| = 2|A| + |B| \leq |A| + |X| = |A| + m$ leading to $|E^+| - m \leq |A|$. On the other hand we have that $|E^+| + |Y_1 \cup Y_2| = 2m$ while $|Y_1 \cup Y_2| + |Y_3| = k - 2 \leq m - 2$. In turn we have that $|Y_3| \leq (m - 2) - |Y_1 \cup Y_2| = (m - 2) - (2m - |E^+|) = |E^+| - m - 2$. However as we have shown earlier, $|Y_3| \geq |A|$, and we have a contradiction. It follows that $f(3, m; 6) \geq 4m + 2$.




Fig. 4.




Fig. 5.

It remains now to construct graphs of girth 6 with degree set $\{3, m\}$ on $4m + 2$ vertices. The approach we use is not unique, it is used as a convenience since it is easy to check that the girth is indeed 6. For $m \geq 5$ consider m 3-remote edges (a_i, b_i) where a_i and b_i are of degree two. Construct a 'ladder' as shown in Fig. 3. Depending on whether or not m is odd or even we add the edges (x, v) and (u, y) or the edges (x, y) and (u, v) . Finally add two new vertices a and b where the neighbours of a and b are $\{a_1, a_2, \dots, a_m\}$ and $\{b_1, b_2, \dots, b_m\}$ respectively. The resulting graphs are shown in Fig. 4i and 4ii. The graphs in Fig. 4 have degree sets $\{3, m\}$ and girth 6 as required. Recalling that $f(3, 3; 6) = 14$, the Heawood graph (see [1]), and that $f(3, 4; 6) = 18$ we have the following.

Theorem 1. *The minimal order of a graph of girth 6 with degree set $\{3, m\}$, $m \geq 3$, is $f(3, m; 6) = 4m + 2$.*

A similar construction using 3-remote vertices instead of edges to form a ladder together with one vertex joined to all of these 3-remote points shows how one may attain the bound of $f(3, m; 5) = 3m + 1$ (see Fig. 5). The case $m = 3$ is the Petersen graph.

The extremal graphs (with respect to number of vertices) are not necessarily unique. For example, if we take a 9-cycle, add three 3-remote vertices of degree three by joining them to every third vertex of this cycle, then we have a 3-regular graph on 12 vertices. Finally join a new vertex to these three extra vertices. The result is a $(3, 4; 5)$ -cage with 21 edges as opposed to a $(3, 4; 5)$ -cage with 20 edges in the earlier construction. This begs the question of how many of these generalized cages there are for a given girth and degree set.

3. The $(4, m; 5)$ -cages

Theorem 2. *The minimal order of a graph of girth 5 with degree set $\{4, m\}$ is $f(4, m; 5) = 4m + 1$, $m \geq 5$.*




Fig. 6.

Proof. Select a vertex x of degree m . Other than including x , the neighbourhoods of the neighbours of x are disjoint (else the girth is 4), thus $f(4, m; 5) \geq 1 + m + 3m = 4m + 1$. It remains then to show that if $m \geq 5$, then we may construct $(4, m; 5)$ -cages of this order.

First define the graph G as follows. Take a $3(m-1)$ -cycle with vertices ordered $0, 1, 2, \dots, 3(m-1)$. Let $X = \{x_1, x_2, \dots, x_{m-1}\}$ and let x_i be adjacent to vertices $(i-1)$, $(m-1) + (i-1)$ and $2(m-1) + (i-1)$ for $i = 0, 1, \dots, m-1$. Define vertices x_{01}, x_{02}, x_{03} where x_{0i} is adjacent to those vertices, j , of C with $j = i-1 \pmod{3}$ for $i = 1, 2, 3$. Let x_0 be a new vertex with neighbours x_{01}, x_{02} and x_{03} and finally join a vertex x to $x_0, x_1, x_2, \dots, x_{m-1}$. Note that in G the vertices x, x_{01}, x_{02} , and x_{03} have degree m while all other vertices are of degree four. It is not difficult to in fact check, for $m \not\equiv 1 \pmod{3}$, that G has the properties we desire. In the rest of our proof however we will only appeal to this construction in the cases $m = 5, 8$ and 9 . Fig. 6 illustrates the case $m = 5$. For all cases other than $m = 5, 8$ or 9 we adapt the construction in Fig. 5-delete one of the long diagonal edges, subdivide two others and finally add two new edges in order to create a graph H with degree set $\{3, m-1\}$, girth 5 and $f(3, m-1; 5) + 2 = 3m$ vertices. The vertices of H are labelled as shown in Fig. 7 where (other




Fig. 7.

than those labelled by m) the labelling is cyclical. H also has the property that the set of vertices labelled i constitute a 3-remote set of vertices for $i = 1, 2, \dots, m$. Now let G be a graph on $4m + 1$ vertices containing a vertex x_0 of degree m with neighbours x_1, x_1, \dots, x_m . The remaining vertices of G induce the subgraph H of Fig. 7. Also, vertex x_i has those vertices labelled i in H as neighbours for $i = 1, 2, \dots, m$. Then G has the desired properties and $f(4, m; 5) = 4m + 1$, $m \geq 5$. \square

When $m = 4$ a different construction leads to the $(4, 5)$ -cage, this is the Robertson graph, (see [1]), the unique $(4, 5)$ -cage with only 19 vertices.

References

- [1] J.A. Bondy and U.S.R. Murty, Graph Theory with Applications (North-Holland, Amsterdam, 1976).
- [2] G. Chartrand, R.J. Gould and S.F. Kapoor, Graphs with prescribed degree sets and girth, Period. Math. Hungar. 12 (1981) 261–266.
- [3] M. M. Downs, R.J. Gould, J. Mitchem and F. Saba, $(D; n)$ -Cages, Congr. Numer. 32 (1981) 279–293.
- [4] M.H. Mulder, Julius Petersen theory of graphs, Institut for Matematik of Datalogi, Odense Universitet, 1990, preprint no. 8.
- [5] P.K. Wong, Cages—A Survey, J. Graph Theory (1982) 1–22.